- 1) A solution is made containing 14.6 g of CH₃OH in 184 g of H₂O. Calculate a) the mole fraction of CH₃OH, b) the mass percent of CH₃OH, and c) the molality of CH₃OH.
- 2) The density of toluene (C_7H_8) is 0.867 g/mL and the density of thiophene (C_4H_4S) is 1.065 g/mL. A solution is made by dissolving 10.0 g of thiophene in 250.0 mL of toluene. Calculate a) the mole fraction of thiophene in the solution, b) the molality of thiophene in the solution, and c) assuming that the volumes of the solute and solvent are additive, what is the molarity of thiophene in solution?
- 3) Calculate the number of moles of solute present in each of the following aqueous solutions: a) 600 mL of 0.250 M SrBr₂, b) 86.4 g of 0.180 m KCl, c) 124.0 g of a solution that is 6.45% glucose ($C_6H_{12}O_6$) by mass.
- 4) Commercial aqueous nitric acid has a density of 1.42 g/mL and is 16 M. Calculate the percent HNO₃ by mass in the solution.
- 5) Commercial concentrated aqueous ammonia is 28% NH $_3$ by mass and has a density of 0.90 g/mL. What is the molarity of this solution?
- 6) Caffeine ($C_8H_{10}N_4O_2$) is a stimulant found in coffee and tea. If a solution of caffeine in chloroform (CHCl₃) as a solvent has a concentration of 0.0750 m, calculate a) the percent caffeine by mass, b) the mole fraction of caffeine.