Unit 6 – Chapter 11	Name
Assignment #2	Period

- 1) A solution is prepared by dissolving 27.0 g of urea, (NH₂)₂CO, in 150.0 g of water. Calculate the boiling point of the solution. Urea is a nonelectrolyte.
- 2) A 2.00-g sample of a large biomolecule was dissolved in 15.0 g of carbon tetrachloride. The boiling point of this solution was determined to be 77.85°C. Calculate the molar mass of the biomolecule. For carbon tetrachloride, the boiling-point constant is 5.03°C ⋅ kg/mol, and the boiling point of pure carbon tetrachloride is 76.50°C.
- 3) The freezing point of *t*-butanol is 25.50° C and K_f is 9.1° C · kg/mol. Usually *t*-butanol absorbs water on exposure to air. If the freezing point of a 10.0-g sample of *t*-butanol is 24.59° C, how many grams of water are present in the sample?
- 4) An aqueous solution of 10.00 g of catalase, an enzyme found in the liver, has a volume of 1.00 L at 27°C. The solution's osmotic pressure at 27°C is found to be 0.74 torr. Calculate the molar mass of catalase.
- 5) A water desalination plant is set up near a salt marsh containing water that is 0.10 *M* NaCl. Calculate the minimum pressure that must be applied at 20.0°C to purify the water by reverse osmosis. Assume NaCl is completely dissociated.
- 6) Calculate the freezing point and the boiling point of each of the following solutions using the observed van't Hoff factors in Table 11.6

a. 0.050 m MgCl₂ b. 0.050 m FeCl₃

7) A 0.500-g sample of a compound is dissolved in enough water to form 100.0mL of solution. This solution has an osmotic pressure of 2.50 atm at 25°C. If each molecule of the solute dissociates into two particles (in this solvent), what is the molar mass of this solute?