


CHEMICAL EQUILIBRIUM B PROBLEMS

Name _____

Period _____


- 1) A mixture of 0.10 moles of NO, 0.050 moles of H₂, and 0.10 moles of H₂O is placed in a 1.0-liter vessel at 300 K. The following equilibrium is established:



At equilibrium, [NO] = 0.062 M.


- Calculate the equilibrium concentrations of H₂, N₂, and H₂O.
- Calculate K_c.

- 2) From the following equation:



Calculate the concentrations of all the species if 0.30 moles of CO and 0.30 moles of O₂ are reacted in a 1.0-liter container.


- 3) A flask is charged with 1.50 atm of N₂O₄ and 1.00 atm NO₂ at 25°C, and the following equilibrium is achieved:



After equilibrium is reached, the partial pressure of NO₂ is 0.512 atm.

- What is the equilibrium partial pressure of N₂O₄?
- Calculate the value of K_p for the reaction.

- 4) At 2000°C, the equilibrium constant for the reaction




If the initial concentration of NO is 0.200 M, what are the equilibrium concentrations on NO, N₂ and O₂?

- 5) For the equilibrium:



At 400 K, $K_c = 7.0$. If 0.30 mol of Br₂ and 0.30 mol Cl₂ are introduced into a 1.0 L container at 400 K, what will be the equilibrium concentrations of Br₂, Cl₂, and BrCl?

- 6) At 218°C, $K_c = 1.2 \times 10^{-4}$ for the equilibrium:



Calculate the equilibrium concentrations of NH₃ and H₂S if a sample of solid NH₄HS is placed in a closed vessel and decomposes until equilibrium is reached.