Name
Period

- 1) A student wants to prepare 1.00 L of a 1.00 M solution of NaOH (molar mass = 40.00 g/mol).
  - a. If solid NaOH is available, how would the student prepare this solution?
  - b. If 2.00 M NaOH is available, how would the student prepare the solution?
  - c. To help insure three significant figures in the NaOH molarity, to how many significant figures should the volumes and mass be determined?
- 2) A solution of ethanol ( $C_2H_5OH$ ) in water is prepared by dissolving 75.0 mL of ethanol (density = 0.79 g/cm<sup>3</sup>) in enough water to make 250.0 mL of solution. What is the **molarity** of the ethanol in this solution?
- 3) If 10.0 g of AgNO₃ is available, what volume of 0.25 M AgNO₃ solution can be prepared?
- 4) How would you prepare 1.00 L of a 0.50 M solution of each of the following?
  - a. H<sub>2</sub>SO<sub>4</sub> from "concentrated" (18 M) sulfuric acid
  - b. HCl from "concentrated" (12 M) reagent
  - c.  $NiCl_2$  from the salt  $NiCl_2 \cdot 6 H_2O$
  - d. HNO<sub>3</sub> from "concentrated" (16 M) reagent
  - e. Sodium carbonate from the pure solid
- 5) Calculate the sodium ion concentration when 70.0 mL of 3.0 *M* sodium carbonate is added to 30.0 mL of 1.0 *M* sodium bicarbonate.
- 6) A stock solution containing Mn<sup>2+</sup> ions was prepared by dissolving 1.584 g pure manganese metal in nitric acid and diluting to a final volume of 1.000 L. The following solutions were then prepared by dilution:

For solution A, 50.00 mL of stock solution was diluted to 1000.0 mL.

For solution B, 10.00 mL of solution A was diluted to 250.0 mL.

For solution C, 10.00 mL of solution B was diluted to 500.0 mL.

Calculate the concentrations of the stock solution and solutions A, B, and C.