Unit 10 – Chapter 5: Gases	Name			
Assignment #4: Practice Problems MC	Period			
1. If a barometer were built using water inst the pressure were 1 atm, knowing that th mercury?	tead of Hg, how high would the column of water be if he density of water is 13.6 times lower than that of			
a. 10.3 m b. 3.17 m	c. 20.0 m d. 33.0 m			
2. Calculate the pressure, in Pascals, for a co a. 2.35 X 10 ⁵ Pa b. 2.70 Pa	olumn of Hg that is 2.05 m high. c. 1.56 X 10 ⁶ Pa d. 2.73 X 10 ⁵ Pa			
3. What is the pressure, in mm Hg, of a gasa. 0.113 mm Hgb. 776 mm Hg	that has a pressure of 15.0 lb/in ² ? c. 1.02 mm Hg d. 27.6 mm Hg			
4. A balloon with an internal pressure of 30 pressure is 15.0 torr. Assuming temperation change?	0.0 torr rises to a height of 30,000 feet, where the ure remains constant, by what ratio did the volume			
a. 25:1 b. 1000:1	c. 20:1 d. 1.85:1			
5. A 0.90-L sample of helium is heated from does this sample occupy at 68°C?	1 68°F to 68°C. At constant pressure, what volume			
a. 1.1 L b. 1.6 L	c. 0.9 L d. 2.7 L			
6. A 3.00-L sample of xenon is heated from increased to 120 cm of Hg. What is the fi	100°F to 200°F and an initial pressure of 70.0 cm nal volume, in L, of the gas?			
a. 1.80 L b. 2.06 L	c. 3.00 L d. 6.00 L			
7. How many moles of an ideal gas are pres 2500 lb/in ² ?	ent in a sample of 1.25 L at 311 K and a pressure of			
a. 8.35 mol b. 5.10 mol	c. 32.8 mol d. 1.02 mol			
 8. A 3.25-L sample of a gas at 80.0°C is heat the final temperature of the gas in Kelvin a. 3.53 X 10³ K b. 151 K 	ed until a final volume of 32.5 L is reached. What is at constant pressure? c. 1.08 X 10 ³ K d. 1.34 X 10 ³ K			
9. Calculate the number of grams of acetyle 20.0°C and a pressure equal to 2500 lb/in	ene, C_2H_2 , in a 30.0 L cylinder at a temperature of ² .			
a. 8.47 X 10 ³ g b. 1000 g	c. 5.52 X 10 ³ g d. 2.40 X 10 ³ g			

__ 10. The average pressure of carbon dioxide in the atmosphere is 25.1 torr at 273 K. Assuming that our atmosphere is 50 miles high and the temperature and pressure of carbon dioxide constant in this region, calculate the mass of the gas in tons. The radius of the earth is 3963.0 miles.

a.	3.0 X 10 ¹⁵ tons	c.	4.8 X 10 ⁶ tons
b.	50,000 tons	d.	6.9 X 10 ⁹ tons

11. A 50.0 L cylinder at temperature of 47°C and a pressure of 50.0 atm contains how many molecules of gas per cm³?

a.	5.73 X 10 ²³	c.	2.30×10^{19}
b.	2.30 X 10 ²²	d.	6.75 X 10 ¹⁸

12. A 50.0 L cylinder of Cl₂ at 20.0°C and a pressure of 103,401 torr springs a leak. The following day the pressure is found to be 41,361 torr. How many moles of chlorine gas escaped during this time?

a.	170 mol	c.	85.0 mol
b.	280 mol	d.	113 mol

13. Tin reacts with hydrochloric acid to produce hydrogen gas and tin (II) chloride. How many liters of hydrogen gas are produced at 27.0° C and a pressure of 710 torr, if 2.80 g of tin reacts with excess hydrochloric acid? Sn_(s) + 2 HCl_(aq) \rightarrow SnCl_{2(aq)} + H_{2(g)} a. 0.620 L c. 2.00 L

a.	0.620 L	с.	2.00 L
b.	0.320 L	d.	1.25 L

14. How many cm³ of carbon tetrachloride are produced when 8.0 L of chlorine are allowed to react with 0.75 L of methane at STP? $4 \operatorname{Cl}_{2(q)} + \operatorname{CH}_{4(q)} \rightarrow 4 \operatorname{HCl}_{(q)} + \operatorname{CCl}_{4(q)}$

			157	(5)	(5)
a.	1500 cm ³	c.	360 cm ³		
b.	750 cm ³	d.	1080 cm	3	

15. Calculate the final pressure, in atm, after 9.06 g of krypton reacts with 10.0 g of fluorine at 300 K in a 10.0 L container. $Kr_{(g)} + F_{2(g)} \rightarrow KrF_{2(s)}$

a. 0.591 atm	c. 0.700 atm
b. 0.384 atm	d. 1.90 atm

16. Calculate the density change, g/L, if 700 g of $C_2H_{6(g)}$ are removed from a 200. 0 L cylinder at 200.0 psi (lb/in²) and a temperature of 20^oC.

a.	3.6 g/L	с.	1.7 g/L
b.	15.0 g/L	d.	16.2 g/L

17. Calculate the density, in g/L, of sulfur dioxide $SO_{2(g)}$ at 37^oC and a pressure of 1440 torr.

a. 6.0 g/L	c. 2.38 g/L
b. 0.60 g/L	d. 4.76 g/L

18. Calculate P_{total}, in atms, for three different gases at partial pressures of 144.0 cm, 800.0 mm, and 1.3 m of Hg.

a.	1.90 atm	c.	1.06 atm
b.	2.58 atm	d.	4.66 atm

19. 1.0 L of hydrogen gas is collected over water at 308 K at a pressure of 728 torr. How many grams of iron are required to react with excess HCl_(aq) to produce this volume of hydrogen gas? The vapor pressure of water is 42.4 torr. The products of the reaction are iron (II) chloride and hydrogen gas.

a.	4.7 g	c.	2.0 g
b.	2.35 g	d.	1.3 g

20. Gas A diffuses twice as fast as gas B. Gas B has a molecular weight = 60.0 g/mol. What is the molar mass of gas A?

a.	15.0 g/mol	с.	30 g/mol
b.	120 g/mol	d.	90 g/mol

_____ 21. If gas B effuses four times as fast as gas A, what is the ratio of the molar masses (A/B)?

a.	2:1	с.	16:1
b.	4:1	d.	8:1

22. The rate of effusion of freon-12 to freon-11 is 1.07:1. The molar mass of freon-11 is 137.4 g/mol. Calculate the molar mass, in g/mol, of freon-12.

a.	100 g/mol	с.	121 g/mol
b.	182 g/mol	d.	118 g/mol

23. Using the van der Waals equation, calculate the pressure exerted by 10.0 g of methane (CH₄) in a 2.1 L container at 330 K. a = 2.253 L³ atm/mol², b = 0.0458 L/mol. Calculate using the ideal gas equation and find the difference between the ideal gas pressure and van der Waals pressure.

a.	2.0 atm	c.	0.1 atm
b.	0.5 atm	d.	1.5 atm